

## A Lax Operator Hierarchy for the New Fifth Order Integrable System

Daryoush Talati\*

Received: 21 July 2015 / Accepted: 4 December 2015

**Abstract** We consider the Lax representation of the new two-component coupled integrable system recently discovered by the author. Connection of the hierarchy of infinitely many Lax pairs with each other is presented.

**Keywords** Symmetry · Lax pair · Integrability

**Mathematics Subject Classification (2010)** 37K15 · 17B80 · 70H06

### 1 Introduction

The term "Lax pair" refers to linear systems that are related to nonlinear equations through a compatibility condition.

The first part of Lax pair is called the scattering problem, that allows the initial-value problem for the integrable equation to be solved exactly. The Lax pair may be used to gather information about the behavior of the solutions to the nonlinear equation.

In his seminal work [2], Lax suggested a formalism to integrate a class of nonlinear evolution equations. He introduced a pair of linear operators  $L$  and  $M$  such that

$$L\phi = \lambda\phi, \quad \phi_t = M\phi, \quad (1)$$

where  $L$  and  $M$  are linear differential operators,  $\lambda$  is an eigenvalue of  $L$ , and  $\phi$  is an eigenfunction of  $L$ . Assuming  $\lambda_t = 0$ , differentiating  $L\phi$  with respect to  $t$  gives

---

\*Corresponding author

Daryoush Talati

Department of Engineering Physics, Ankara University 06100 Tandoğan, Ankara, Turkey

E-mail: Daryoush.talati@eng.ankara.edu.tr

$$(L_t + [L, M])\phi = 0. \quad (2)$$

Where  $[M, L] = ML - LM$  is the operator commutator. Hence,

$$L_t = [M, L], \quad (3)$$

is called Lax equation and contains commutative nonlinear evolution equation for suitable  $L$  and  $M$ . In [1], it is shown that all scalar evolution equations, which have the form of a conservation law, have a Lax pair with a second order  $L$ . For example consider the Lax formalism for the KdV equation

$$L = -\mathcal{D}_x^2 + u + u_1 \mathcal{D}_x^{-1}, \quad M = -4\mathcal{D}_x^3 + 6u\mathcal{D}_x + 9u_1 + 3u_2 \mathcal{D}_x^{-1}. \quad (4)$$

These operators satisfies the Lax equation (3 if  $u$  is solution to the KdV equation:

$$u_t = -u_3 + 6uu_1. \quad (5)$$

Very recently, a integrable system of fifth order nonlinear equations founded by the author [3]

$$\begin{pmatrix} u \\ v \end{pmatrix}_t = \begin{pmatrix} u_{xxxxx} - vv_{xxxx} + 10u_x u_{xxx} - 20u^2 u_{xxx} \\ -4v_x v_{xxx} - 2uv_{xxx} + 10u_x^2 - 80u u_x u_{xx} \\ -32v v_x u_{xx} - 3v_x^2 - 18v u_x v_{xx} - 6w v_x v_{xx} \\ +16u^2 v v_{xx} + 8v^3 v_{xx} - 20u_x^3 - 18u_x v_x^2 \\ +32uv u_x v_x + 80u^4 u_x + 48u^2 v^2 u_x + 4v^4 u_x \\ +16u^2 v_x^2 + 24v^2 v_x^2 + 32u^3 v v_x + 16uv^3 v_x \\ -8v^2 u_{xxx} \\ 2v u_{xxxx} + 2v_x u_{xxx} - 4uv u_{xxx} - 2v^2 v_{xxx} \\ -4w v_x u_{xx} - 32u^2 v u_{xx} - 16v^3 u_{xx} - 8v v_x v_{xx} \\ +12u_x^2 v_x - 64uv u_x^2 - 48v^2 u_x v_x + 64u^3 v u_x \\ -32u^2 u_x v_x + 32uv^3 u_x - 2v_x^3 + 16u^4 v_x \\ +48u^2 v^2 v_x + 20v^4 v_x + 20v u_x u_{xx} \end{pmatrix}. \quad (6)$$

Talati [3] proved that this system is bi-Hamiltonian and therefore possess infinitely many generalized symmetries. Almost all of the known integrable models possess linear Lax pairs. However in the literature, there is no systematic way of finding whether a given evolution equation possesses a Lax representation.

## 2 Lax pair of system (6)

The aim of this paper is to construct a Lax pair for the new fifth order integrable system introduced by the author [3]. For the system (6) we shall consider the case where our candidate Lax pair is a differential operator

$$\begin{aligned}\mathcal{M} &= \alpha_0 \mathcal{D}_x^i + \alpha_1 \mathcal{D}_x^{i-1} + \dots + \alpha_{i-1} \mathcal{D}_x + \alpha_i, \\ \mathcal{L} &= \beta_0 \mathcal{D}_x^j + \beta_1 \mathcal{D}_x^{j-1} + \dots + \beta_{j-1} \mathcal{D}_x + \beta_j.\end{aligned}\tag{7}$$

Here  $\alpha, \beta = \alpha, \beta(u, v, u_x, v_x, \dots)$  are the dependent variables and  $i, j = 1, 2, 3, \dots$ . Now the major problem is to find appropriate analytic functions  $\alpha_i$  and  $\beta_j$  that satisfy the integrability condition (3). From the order of the system (6) it is easy to see that we must assign  $i = 3$

$$\mathcal{M} = \alpha_0 \mathcal{D}_x^5 + \alpha_1 \mathcal{D}_x^4 + \alpha_2 \mathcal{D}_x^3 + \alpha_3 \mathcal{D}_x^2 + \alpha_4 \mathcal{D}_x + \alpha_5.\tag{8}$$

By assuming that  $\alpha_0$  and  $\beta_0$  are nonzero constants, We carry out the calculations For cases  $j = 1, 2, 3, \dots$  according to (3). After lengthy calculations, the integrability condition (3) can lead to the following Lax pair:

$$\begin{aligned}\alpha_1 &= \frac{1}{2}(10u + 5v), \\ \alpha_2 &= \frac{1}{2}(20u_x + 10v_x + 20u^2 + 20uv + 5v^2), \\ \alpha_3 &= \frac{1}{4}(40u_{xx} + 20v_{xx} + 120u_x u + 60u_x v + 60v_x u + 30v_x v + 40u^3 \\ &\quad + 60u^2 v + 30uv^2 + 5v^3), \\ \alpha_4 &= \frac{1}{16}(80u_{xxx} + 40v_{xxx} + 320u_{xx} u + 160u_{xx} v + 160v_{xx} u + 80v_{xx} v \\ &\quad + 240u_x^2 + 240u_x v_x + 480u_x u^2 + 480u_x uv + 120u_x v^2 + 60v_x^2 \\ &\quad + 240v_x u^2 + 240v_x uv + 60v_x v^2 + 80u^4 + 160u^3 v + 120u^2 v^2 \\ &\quad + 40uv^3 + 5v^4), \\ \alpha_5 &= \frac{1}{32}(16v_{xxxx} + 160u_{xxx} u + 48u_{xxx} v + 80v_{xxx} u + 72v_{xxx} v + 160u_{xx} v_x \\ &\quad + 960u_{xx} u^2 + 384u_{xx} uv + 336u_{xx} v^2 + 160v_{xx} u_x + 176v_{xx} v_x \\ &\quad + 160v_{xx} u^2 + 224v_{xx} uv + 72v_{xx} v^2 + 1120u_x^2 u + 48u_x^2 v \\ &\quad + 480u_x v_x u + 752u_x v_x v + 320u_x u^3 + 992u_x u^2 v + 240u_x uv^2 \\ &\quad + 296u_x v^3 + 184v_x^2 u + 92v_x^2 v + 160v_x u^3 - 272v_x u^2 v \\ &\quad + 120v_x uv^2 - 236v_x v^3 - 480u^5 - 176u^4 v - 432u^3 v^2 \\ &\quad - 216u^2 v^3 - 118uv^4 - 63v^5), \\ \mathcal{L}_{j=1} &= \frac{1}{2}D_x + \frac{u}{2} + \frac{v}{4}, \\ \mathcal{L}_{j=2} &= D_x^2 + (2u + v)D_x + \frac{1}{4}(4u_x + 2v_x + 4u^2 + 4uv + v^2).\end{aligned}\tag{9}$$

Using the previous result we will state a conjecture about the existence a hierarchy of infinitely many Lax operators of the system (6).

**Conjecture 1:** Consider the differential operators

$$\mathcal{L}_j = \sum_{n=0}^j \beta_n D_x^{j-n}, \quad j = 1, 2, 3, \dots \quad (10)$$

Where

$$\beta_n = \frac{j-n+1}{n} \left( \frac{d}{dx} + u + \frac{1}{2}v \right) \beta_{n-1}, \quad \beta_0 = \frac{j}{2}. \quad (11)$$

Then it can be further proved that the operator equation  $\mathcal{L}_{j_t} - [\mathcal{M}, \mathcal{L}_j] = \mathcal{O}$  where  $\mathcal{O}$  is the zero operator, is equivalent to the system (6) in the sense that both sides of the equation turn out to be operators of multiplication by a function. So the system (6) admits a hierarchy of infinitely many Lax operators that satisfy the condition  $[\mathcal{L}_n, \mathcal{L}_m] = 0$ .

## References

1. F. Calogero and M.C. Nucci, Lax pairs galore, *J. Math. Phys.*, **32**, 72-74 (1991).
2. P.D. Lax, Integrals of nonlinear equations of evolution and solitary waves, *Comm. Pure Appl. Math.*, **21**, 467-490 (1968).
3. D. Talati, A fifth-order bi-Hamiltonian system. arXiv preprint arXiv:1304.1987 (2013).