

Eccentric Connectivity Index of Nanostar Dendrimer $NS_3[n]$

Morteza Alishahi

Received: 29 November 2023 / Accepted: 20 April 2024

Abstract Let G be a molecular graph. The eccentric connectivity index, $\zeta^c(G)$, is defined as, $\zeta^c(G) = \sum_{u \in V(G)} deg(u)ecc(u)$, where $deg(u)$ denotes the degree of vertex u and $ecc(u)$ is the largest distance between u and any other vertex v of G . In this paper, an exact formula for the eccentric connectivity index of nanostar dendrimer $NS_3[n]$ is given.

Keywords Eccentric connectivity index · Nanostar dendrimer · Topological index.

1 Introduction

Chemical graph theory is one of the branches of mathematical chemistry. In chemical graph theory, a variety of concepts from graph theory are used to model chemical phenomena graphically. In this modeling, each atom is represented by a vertex and each bond between two atoms is represented by an edge. A topological index for an undirected simple graph G is a numerical value that is invariant under all graph isomorphisms which correlates to its Physico-chemical properties. Topological indices are used for studying QSAR (quantitative structure-activity relationships) and QSPR (quantitative structure-property relationships) for foretelling many attributes of chemical compounds and their biological properties. Various studies have been performed on different topological indices [1–11].

Dendrimers are highly ordered, branched polymeric molecules. They have many applications in gene therapy, nanotechnology, medicine production, and other fields. Every dendrimer is a macromolecule which made of a core with

M. Alishahi
Department of Mathematics, Nazarabad Center, Karaj Branch, Islamic Azad University,
Karaj, Iran.
E-mail: morteza.alishahi@gmail.com

tree-like arms or branches named dendrons. The zero generation of a dendrimer is the core molecule of dendrimer without dendrons. Each generation of a dendrimer is made by adding some new branches along the branches of the previous generation with a specific rule. Our aim in this study is to investigate a special topological property of dendrimers. The molecular graph of a molecule M is a graph with the finite set of all atoms as its vertex set and chemical bonds are the edges of this graph. We use the notations $G(M)$, G for short, for this graph, $V(G)$ for its vertex set, and $E(G)$ for the set of all edges. For each vertex u , $\deg(u)$ denotes the degree of u . If $x, y \in V(G)$, then the length of a minimum path connecting x and y is named the distance between x and y and denoted by $d(x,y)$.

Sharma, Goswami, and Madan proposed the eccentric connectivity index of the molecular graph G which is defined as

$$\zeta^c(G) = \sum_{u \in V(G)} \deg(u) \text{ecc}(u),$$

where $\text{ecc}(u) = \max\{d(u,v) | v \in V(G)\}$ [11].

Ashrafi and Saheli computed the eccentric connectivity index of nanostar dendrimers $NS_1[n]$ and $NS_2[n]$, see [3,10] for details. In this study, we are going to compute the eccentric connectivity index of nanostar dendrimer $NS_3[n]$.

2 MAIN RESULTS AND DISCUSSION

$NS_1[n]$, $NS_2[n]$ and $NS_3[n]$ are three types of dendrimers with n generations. $NS_1[n]$ (for $n=3$) is depicted in Fig. 1 and its generator is shown in Fig. 2.




Fig. 1 The molecular graph of $NS_1[3]$

In [10], Saheli and Ashrafi computed the eccentric connectivity index of nanostar dendrimer $NS_1[n]$ as

$$\zeta^c(NS_1[n]) = 135n \times 2^{n+2} + 135 \times 2^n - 50n + 179.$$




Fig. 2 The core of $NS_1[n]$

$NS_1[n]$ (for $n=2$), and its core, are shown in Figs. 3 and 4, respectively. In [3], Ashrafi and Saheli computed the eccentric connectivity index of $NS_2[n]$ as

$$\zeta^c(NS_2[n]) = 420n \times 2^n + 60 \times 2^n - 110n + 40.$$




Fig. 3 The molecular graph of $NS_2[2]$




Fig. 4 The core of $NS_2[n]$

Now, we consider nanostar dendrimer where (Figs. 5 and 6). In the following we try to compute the eccentric connectivity index of $NS_3[n]$.

Theorem 1 *The eccentric connectivity index of nanostar dendrimer $NS_3[n]$, is computed as*

$$\zeta^c(NS_3[n]) = 29n \times 2^{n+5} + 741 \times 2^{n+1} - 104n - 98, n \geq 1.$$

Proof Considering Figs. 7, 8 and Table 1. It can be seen that there exist 22 types of vertices in $NS_3[n]$, based on their positions in branches of $NS_3[n]$




Fig. 5 The molecular graph of $NS_3[1]$




Fig. 6 The molecular graph of $NS_3[2]$

(Fig. 8). Therefore, we have:

$$\begin{aligned}
 \zeta^c(NS_3[n]) &= \sum_{u \in V(NS_3[n])} \deg(u) \text{ecc}(u) \\
 &= 2 \times (8n + 19) \times 2^{n+2} + 2 \times (8n + 18) \times 2^{2n+2} + 3 \times (8n + 17) \times 2^{n+1} \\
 &\quad + 2 \times (8n + 16) \times 2^{n+1} + 2 \times (8n + 15) \times 2^{n+1} + 3 \times (8n + 14) \times 2^{2n+1} \\
 &\quad + 2 \times (8n + 15) \times 2^{n+1} + 3 \times (8n + 13) \times 2^{n+1} + 3 \times (8n + 18) \times 2^{n+1} \\
 &\quad + 2 \times (8n + 17) \times 2^{2n+1} + 2 \times (8n + 16) \times 2^{n+1} + 3 \times (8n + 15) \times 2^{n+1} \\
 &\quad + 2 \times (8n + 16) \times 2^{n+1} + 3 \times (8n + 14) \times 2^{2n+1} + 1 \times (8n + 15) \times 2^{n+1} \\
 &\quad + 2 \times (8n + 12) \times 2^{n+1} + 2 \times (8n + 11) \times 2^{n+1} \\
 &\quad + 2 \times \sum_{k=0}^{n-1} (8n - 4k + 10) \times (2^{n-k+1}) + 2 \times \sum_{k=0}^{n-1} (8n - 4k + 9) \times (2^{n-k+1}) \\
 &\quad + 2 \times (8n + 16) \times 2^{n+1} + 2 \times \sum_{k=0}^{n-1} (8n - 4k + 8) \times (2^{n-k+1}) \\
 &\quad + 3 \times \sum_{k=0}^{n-2} (8n - 4k + 7) \times (2^{n-k}) + 3 \times (4n + 11) \times 2 + 2 \times (4n + 10) \times 2 \\
 &= 29n \times 2^{n+5} + 741 \times 2^{n+1} - 104n - 98
 \end{aligned}$$

Table 1 Types of vertices in $NS_3[n]$.

Layer	Types of vertices	Frequencies	$\text{ecc}(u)$	$\text{deg}(u)$
n	1	2^{n+2}	$8n+19$	2
n	2	2^{n+2}	$8n+18$	2
n	3	2^{n+1}	$8n+17$	3
n	4	2^{n+1}	$8n+16$	2
n	5	2^{n+1}	$8n+15$	2
n	6	2^{n+1}	$8n+14$	3
n	7	2^{n+1}	$8n+13$	3
n	8	2^{n+1}	$8n+15$	2
n	9	2^{n+1}	$8n+18$	3
n	10	2^{n+1}	$8n+17$	2
n	11	2^{n+1}	$8n+16$	2
n	12	2^{n+1}	$8n+15$	3
n	13	2^{n+1}	$8n+16$	2
n	14	2^{n+1}	$8n+14$	3
n	15	2^{n+1}	$8n+15$	1
n	16	2^{n+1}	$8n+12$	2
n	17	2^{n+1}	$8n+11$	2
n	18	2^{n+1}	$8n+10$	2
n	19	2^{n+1}	$8n+9$	2
n	20	2^{n+1}	$8n+8$	2
n	21	2^n	$8n+8$	3
For $1 \leq i \leq n-1$				
i	18	2^{i+1}	$8i+14$	2
i	19	2^{i+1}	$8i+13$	2
i	20	2^{i+1}	$8i+12$	2
i	21	2^i	$8i+11$	3
i	22	2	$4i+10$	2




Fig. 7 The eccentricity of vertices in a quarter of $NS_3[2]$




Fig. 8 Types of vertices in a quarter of $NS_3[2]$

References

1. A. R. Ashrafi, A. Loghman, PI index of armchair polyhex nanotubes, *Ars Comb.*, 80, 193–199 (2006).
2. A. R. Ashrafi, M. Mirzargar, PI, Szeged, edge Szeged indices of an infinite family of nanostar dendrimers, *Indian J. Chem.*, 47A, 538–541 (2008).
3. A. R. Ashrafi, M. Saheli, Computing eccentric connectivity index of a class of nanostar dendrimers, *Kragujevac J. Sci.*, 34, 65–70 (2012).
4. N. De, S. M. A. Nayeem, A. Pal, Computing modified eccentric connectivity index and connective eccentric index of V-phenylenic nano torus, *Stud. UBBChem.*, 59, 129–137 (2014).
5. I. Gutman, P. Khadikar, P. Rajput, S. Karmarkar, The Szeged index of polyacenes, *J. Serb. Chem. Soc.*, 60, 759–764 (1995).
6. I. Gutman, O. E. Polansky, *Mathematical Concepts in Organic Chemistry*, Springer-Verlag, New York (1986).
7. A. Hedyari, B. Taeri, Wiener and Schultz indices of TUC4C8(R) nanotubes, *J. Comp. Thoe. Nanosci.*, 4, 158–167 (2007).
8. A. Iranmanesh, N. A. Gholami, Computing the Szeged index of styryl benzene dendrimer and triarylamine dendrimer of generation 1-3, *Math. Comput. Chem.*, 62, 371–379 (2009).
9. A. Karbasioun, A. R. Ashrafi, Wiener and detour indices of a new type of nanostar dendrimers, *Macedonian J. Chem. Eng.*, 28, 49–54 (2009).
10. M. Saheli, A. R. Ashrafi, The eccentric connectivity index of nanostar dendrimers, *International Journal of Chemical Modeling.*, 3, 227–232 (2011).
11. V. Sharma, R. Goswami, A. K. Madan, Eccentric connectivity index: A novel highly discriminating topological descriptor for structure-property and structure-activity studies, *J. Chem. Inf. Comput. Sci.*, 37, 273–282 (1997).